

STARTING THE CONVERSATION

Only you and your healthcare provider know what treatment is right for you. Here are some questions you can use to start a conversation with your healthcare provider about your treatment options.

- How will my treatment fit into my routine?
- What side effects could I experience?
- Can I still take other medications with my HIV-1 medication?
- Do I need to eat when I take my HIV-1 medication?
- Are there other health risks my HIV or HIV-1 treatment could cause?



YOUR HEALTH AND HIV-1

Staying undetectable

PIFELTRO, when taken as part of a multi-tablet regimen, can help you get your viral load to undetectable. **Undetectable means the level of HIV-1 in your blood is so low that lab tests can't detect it.** And when your HIV-1 is under control, you can focus on your health and life goals beyond HIV-1.

What is PIFELTRO?

PIFELTRO is a prescription HIV medicine used with other antiretroviral medicines to treat human immunodeficiency virus-1 (HIV-1) infection in adults:

- who have not received HIV-1 medicines in the past,
- or to replace their current HIV-1 medicines for people whose healthcare provider determines that they meet certain requirements.

HIV-1 is the virus that causes AIDS (Acquired Immune Deficiency Syndrome).

Please read the [Patient Information](#) for PIFELTRO and discuss it with your healthcare provider. The physician [Prescribing Information](#) also is available.

Important Safety Information

Do not take PIFELTRO if you are currently taking any of the following medicines:

- | | |
|-----------------|-------------------|
| • carbamazepine | • rifampin |
| • oxcarbazepine | • rifapentine |
| • phenobarbital | • mitotane |
| • phenytoin | • St. John's wort |
| • enzalutamide | |

Ask your healthcare provider or pharmacist if you are not sure if your medicine is one that is listed above. If you have taken any of the medicines in the past 4 weeks, talk to your healthcare provider or pharmacist before starting PIFELTRO.

Important Safety Information continued on next page.



Important Safety Information (continued)

PIFELTRO can cause serious side effects, including:

Severe skin reactions have happened in people treated with PIFELTRO. Call your healthcare provider right away if you develop a rash during treatment with PIFELTRO. **Stop taking PIFELTRO and get medical help right away if you develop a painful rash with any of the following symptoms:** fever, blisters or sores in the mouth, blisters or peeling of the skin, or redness or swelling of the eyes (conjunctivitis).

Changes in your immune system (Immune Reconstitution Syndrome) can happen when you start taking HIV-1 medicines. Your immune system may get stronger and begin to fight infections that have been hidden in your body for a long time. Tell your healthcare provider right away if you start having new symptoms after starting your HIV-1 medicine.

The most common side effects of PIFELTRO include: nausea, dizziness, headache, tiredness, diarrhea, stomach (abdominal) pain, and abnormal dreams.

These are not all the possible side effects of PIFELTRO. For more information, ask your healthcare provider or pharmacist.

Before starting PIFELTRO, tell your healthcare provider about all your medical conditions, including if you are pregnant or plan to become pregnant. It is not known if PIFELTRO can harm your unborn baby. Tell your healthcare provider if you are breastfeeding or plan to breastfeed. PIFELTRO may pass to your baby in your breast milk. HIV may pass to your baby through your breast milk if your baby does not have HIV.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. **Some medicines interact with PIFELTRO.** Keep a list of your medicines to show your healthcare provider and pharmacist. Tell your healthcare provider if you have taken rifabutin in the past 4 weeks. **Do not start taking a new medicine without telling your healthcare provider.** Your healthcare provider can tell you if it is safe to take PIFELTRO with those other medicines.

Take PIFELTRO every day exactly as your healthcare provider tells you to take it. Take PIFELTRO 1 time each day, at about the same time every day.

Tell your healthcare provider if you have taken rifabutin in the past 4 weeks.

If you take the medicine rifabutin during treatment with PIFELTRO, take PIFELTRO 2 times each day, about 12 hours apart, as prescribed by your healthcare provider. You may not have enough doravirine in your blood if you take rifabutin during treatment with PIFELTRO.

Do not change your dose or stop taking PIFELTRO without talking to your healthcare provider. Stay under a healthcare provider's care when taking PIFELTRO.

Take PIFELTRO on a regular dosing schedule as instructed by your healthcare provider. Do not miss doses.

If you miss a dose of PIFELTRO, take it as soon as you remember. If it is almost time for your next dose, skip the missed dose and take the next dose at your regular time. Do not take 2 doses of PIFELTRO at the same time.

Get your PIFELTRO refilled from your healthcare provider or pharmacy before you run out.

When your PIFELTRO supply starts to run low, get more from your healthcare provider or pharmacy. This is very important because the amount of virus in your blood may increase if the medicine is stopped for even a short time. The virus may develop resistance to PIFELTRO and become harder to treat.

Please read the [Patient Information](#) for PIFELTRO and discuss it with your healthcare provider. The physician [Prescribing Information](#) also is available.



Copyright © 2025 Merck & Co., Inc., Rahway, NJ, USA and its affiliates.
All rights reserved. US-DOV-01593 01/25 www.pifeltro.com

Pifeltro[®]
doravirine
100 mg tablets